

SUPERCONDUCTIVITY OF NEARLY 2-D SYSTEMS

by

Nguyen Minh Khue

Institute of Theoretical Physics

National Centre for Natural Science and Technology of Vietnam

P.O.Box 429, Hanoi 10000, Vietnam

and

Le Ngoc Minh

Hue University, Hue City, Vietnam

and

Nguyen Tri Lan

Institute of Theoretical Physics

National Centre for Natural Science and Technology of Vietnam

P.O.Box 429, Hanoi 10000, Vietnam

and

Do Huu Nha

National Pedagogic Institute of Vietnam

Dich vong, Tu liem, Hanoi, Vietnam

PACS numbers: 74.20.-z

Abstract

Effect of geometry on the superconductivity is considered. It is shown that the for nearly two dimensional BCS systems the critical temperature is rapidly increased with decreasing the thickness of the layer. The result is expected to be useful in the explanation of high-Tc conductivity of the ceramic compounds.

1. Introduction

In recent years finding mechanisms for high-Tc superconductivity is one of the most interesting topic in Condensed matter Physics. Commonly it is believed that the reasons leading to the high-Tc superconductivity of ceramic compounds are the strong correlation between electrons or the two-dimensional (2-D) character of the crystals or both of them. In almost theories ideal 2-D systems have been considered [1]. In this work we consider the superconductivity of nearly 2-D BCS systems and show that for such systems the critical temperature T_c is very sensitive to the thickness d of the layer, Relation between T_c and d is established in section 3 basing on the model described in Section 2.

2. The Model

Let us consider a nearly 2-D interacting electron system with thickness d . In the coordinate system with z-axis perpendicular to the layer the system can be described by the following Hamiltonian

$$H = \sum_i \left[-\frac{\hbar^2}{2m} \nabla_i^2 + v(\vec{r}_i) \right] + \frac{1}{2} \sum_{i \neq j} V(\vec{r}_i - \vec{r}_j) \quad (1)$$

where

$$v(\vec{r}) = \begin{cases} 0 & \forall x, y \quad \text{and} \quad -d/2 \leq z \leq d/2 \\ \infty & \forall x, y \quad \text{and} \quad z < -d/2 \quad \text{or} \quad z > d/2 \end{cases} \quad (2)$$

The first term in the r.h.s. of (1) describes non-interacting electrons, its eigenfunctions and eigenvalues are

$$\varphi_{n\vec{K}} = (dS)^{-1/2} \sin\left(\frac{\pi n z}{d}\right) e^{i\vec{K}\vec{X}} \quad (3)$$

$$\epsilon_{n\vec{K}} = \frac{(\pi\hbar n)^2}{2md^2} + \frac{\hbar^2 \vec{K}^2}{2m}, \quad n = 1, 2, 3... \quad (4)$$

in which $\vec{K}$ and $\vec{X}$ are 2-D vectors : $\vec{K} = (k_x, k_y)$, $\vec{X} = (x, y)$, and S is the area of the system. In the following the symbols $\epsilon_n \equiv (\pi\hbar n)^2/2md^2$, $\epsilon_{\vec{K}} \equiv \hbar^2 \vec{K}^2/2m$ will be used and $\epsilon_{\vec{K}}$ will be measured from $\epsilon_{\vec{K}_F}$ with $\vec{K}_F$ being the Fermi momentum. The structure of the spectrum is illustrated in Fig.1. The occupation of single states depends on the thickness, some possibilities are shown in Fig.2.

Using (3) as basic functions for the second quantization and remaining only matrix elements describing the scattering between electrons of opposite momentum and spin we have

$$H = \sum_{nK} \epsilon_{nK} a_{nK}^+ a_{nK} + \frac{1}{2} \sum_{nK, n'K'} V_{nK, n'K'} a_{n,K}^+ a_{n,-K}^+ a_{n',-K'} a_{n',K'} \quad (5)$$

in which $a_{n,K}^+$ and $a_{n,K}$ are the creation and annihilation operators for electron on branch n with momentum $\vec{K}$ and spin up and $a_{n,-K}^+$ and $a_{n,-K}$ are that for electron on branch n with momentum $-\vec{K}$ and spin down.

As in BCS theory the sum in the second term in the r.h.s. of (5) will be restricted by the condition $|\epsilon_{\vec{K}}| \leq \hbar\omega_D$ with ω_D being the Debye frequency.

3. Superconductivity

Following BCS we consider the pairing between electrons with opposite momentum and spin with energy $|\epsilon_{\vec{K}}| \leq \hbar\omega_D$, i.e, we chose the trial function in the following form

$$|\psi\rangle = \prod_{n,K} (u_{n,K} + v_{n,K} a_{n,K}^+ a_{n,-K}^+) |O\rangle \quad (6)$$

with the conditions

$$u_{n,K}^2 + v_{n,K}^2 = 1 \quad (7)$$

$$u_{n,-K} = u_{n,K}, \quad v_{n,-K} = -v_{n,K} \quad (8)$$

Note that in state $|\psi\rangle$ there are only pairs of electrons on the same branch of spectrum since the two conditions $|\epsilon_{\vec{K}}| \leq \hbar\omega_D$ and $|\epsilon_{-\vec{K}}| \leq \hbar\omega_D$ can not be satisfied simultaneously for different branches.

Carrying the same calculations as in the traditional BCS theory [2] we obtain the following system of equations for the gaps

$$\Delta_{n,K} = -\frac{1}{2} \sum_{n',K'} V_{nK, n'K'} \frac{\Delta_{n',K'}}{E_{n',K'}} \tanh \frac{1}{2} \beta E_{n',K'} \quad (9)$$

in which $E_{nK} = (\epsilon_{nK}^2 + \Delta_{nK}^2)^{1/2}$, $\beta = 1/K_B T$, K_B is the Boltzman constant and T is the temperature.

After investigating Eq.(9) we note that for small d the interaction $V(\vec{r}_i - \vec{r}_j)$ is weakly dependent on $z_i - z_j$ therefore we may assume that $V(\vec{r}_i - \vec{r}_j) = V(\vec{X}_i - \vec{X}_j)$ and therefore $V_{n,K,n'K'} = \delta_{nn'} V_{K,K'}$. We consider two cases.

(i) For the case shown in Fig.2a we may assume

$$V_{K,K'} = \begin{cases} -(V/S) & \text{if } |\epsilon_K| \text{ and } |\epsilon_{K'}| \leq \hbar\omega_D \text{ and spins in states } K \text{ and } K' \\ & \text{are the same and } K \text{ and } K' \text{ belong to branch } n = 1 \\ 0 & \text{otherwise} \end{cases} \quad (10)$$

Substituting (10) into Eq.(9) we obtain the following equation for the temperature β_{1c} for which $\Delta_1(\beta_{1c}) = 0$.

$$\int_0^{\hbar\omega_D} \frac{d\xi}{\xi + \epsilon_1} th \frac{1}{2} \beta_{1c} (\xi + \epsilon_1) = \frac{1}{N(0)V} \quad (11)$$

in which $N(0)$ is the density of states corresponding to $\epsilon_{\vec{K}}$ at the Fermi level.

(ii) For the case shown in Fig.2b we may assume

$$V_{K,K'} = \begin{cases} -(V/S) & \text{if } |\epsilon_K| \text{ and } |\epsilon_{K'}| \leq \hbar\omega_D \text{ and spins in states } K \text{ and } K' \\ & \text{are the same and both } K \text{ and } K' \text{ belong to branch } n = 1 \text{ or } n = 2 \\ 0 & \text{otherwise} \end{cases} \quad (12)$$

Substituting (12) into (9) we see $\Delta_{nK} = 0$ for $n \neq 1, 2$ and $\Delta_{nK} = 0$ if $|\epsilon_K| > \hbar\omega_D$, $\Delta_{nK} = \Delta_n$ if $|\epsilon_K| \leq \hbar\omega_D$ for $n = 1, 2$ and Δ_n is the solution of the following equation

$$\left[1 - \frac{V}{2S} \sum_K \frac{1}{E_{nK}} th \frac{1}{2} \beta E_{nK}\right] \Delta_n = 0. \quad (13)$$

The nontrivial solution of Eqs.(13) can be found by solving the following equation.

$$\frac{V}{2S} \sum_K \frac{1}{E_{nK}} th \frac{1}{2} \beta E_{nK} = 1. \quad (14)$$

Setting $\Delta_n(\beta_{nc}) = 0$ ($n = 1, 2$) in (14) we obtain the following equation

$$\int_0^{\hbar\omega_D} \frac{d\xi}{\xi + \epsilon_n} th \frac{1}{2} \beta_{nc} (\xi + \epsilon_n) = \frac{1}{N(0)V}, \quad (n = 1, 2) \quad (15)$$

Eqs.(11) and (15) give us

$$K_B T_{nc} = 1, 14(\hbar\omega_D + (n\pi\hbar)^2/2md^2) e^{-1/N(0)V}, \quad (n = 1, 2) \quad (16)$$

By the definition the critical temperature $T_c = T_{1c}$ since $T_{1c} < T_{2c}$

Formula (16) shows that the critical temperature is increased with decreasing the thickness and we hope that it should be useful in the explanation of the high- T_c superconductivity of the ceramic compounds.

ACKNOWLEDGEMENTS

One of the authors (NMK) should like to express his deep gratitude to Prof. A.Zawadowski, Prof. J.Sólyom, Dr. P.Fazekas and Dr. F.Woynarovich for permanent paying attention to him.

The work has been completed with the financial support from Vietnam National Research Program in Natural Sciences.

REFERENCES

1. A.P. Balachandran, E.Ercolessi, G.Morandi and Ajit Mohan Srivastava., *The Hubbard Model and Anyon Superconductivity*, SU-4228-431, TPI-MINN-90/26-T May 1990.
2. J.R.Schrieffer, *Theory of Superconductivity*, Benjamin, 1964.

FIGURE CAPTIONS

Fig.1. The structure of the spectrum.

Fig.2. Some possibilities of the occupation of single particle states.